

## Vaccine damage and its repercussions – Diagnosis and therapy with BICOM

Anita Klammrodt, Naturopath, Grossefehn

### INTRODUCTION

My name is Anita Klammrodt. I have been a naturopath for eleven years and have been using bioresonance for diagnosis and treatment in my practice for seven years. I use bioresonance among other things to treat the side effects and damage caused by vaccines.

I first became aware of this through my son. He was given a 5-way vaccine at the age of three when he was due to go to nursery school. As a result of this he changed markedly. From being a loving, happy little boy he turned into an unsettled, unhappy child with destructive tendencies. Naturally, I did not think that the multiple vaccine could be the cause. I simply believed that he had picked up his bad behaviour from other children in the nursery.

In 1985, when our son was almost 15 years old, we joined a self-help group for hyperactive children which initially focused mainly on nutrition. However, attention soon turned to inoculations too. Speaking to other parents in the group, the subject of inoculations kept coming up. Some parents had noticed that following inoculations their children had become more hyperactive, cried all the time or developed neurodermatitis.

### PATIENTS WITH ADS, LACK OF CONCENTRATION, MOTOR-RELATED PROBLEMS, PROBLEMS AT SCHOOL

Now, since becoming a naturopath, a number of parents come to me with their children who are suffering from ADS, poor concentration, motor-related problems and who do badly in school. These children, despite having a normal level of intelligence, experience difficulties at school, their handwriting is very bad and they get bad marks for their work despite intensive practice at home. Some wet their beds until they are 13 years old.

Their short-term memory is very poor. Some children also develop nervous ticks. Their development is often severely delayed and their motor activity is disturbed. Since I have been working with the bioresonance method I have established that children with behavioural difficulties very often suffer from an inoculation stress and can be helped effectively with BICOM.

I test all stresses with program 170. At this point I would like to report on three cases which are typical of those occurring at my practice.

### Case 1: ADS and learning difficulties

A boy, aged 10, has enormous problems with concentration at school and teachers complain that he is often disruptive. He never wants to do what the teacher tells him. He has already repeated a year of school and just doesn't go anymore. His mother explained that her child often "can't think straight". Despite their best efforts homework is a major challenge for both mother and child.

#### Testing

Using the BICOM I established that this child displayed a strong stress through the tetanus-diphtheria-Hib-whooping cough-polio inoculation.

#### Therapy

Program: 191 detoxification

Time: 6 mins.

Amplification: 6

IC: immune serum + blood

Program: 930 lymph activation  
430 liver detoxification

IC: blood

I: flexible electrode on stomach and forehead

O: modulation mat

MD: chip

Treated once every week.

After 6 treatments the mother reported that her child was now completing his homework much more quickly and the teachers were also saying that her son was able to learn more effectively, was more attentive and no longer disruptive in lessons.

## Case 2: Problems with concentration

Boy, 13 years old. Mother reported that since his tetanus booster he had been getting bad grades at school and could no longer concentrate. He was always forgetting his school things. The boy told me himself when his mother was not there that he felt miserable all the time and preferred to stay at home in his room. He also felt a constant pressure in his head.

### Testing

My testing for vaccine damage was positive.

### Therapy

Program: 191 detoxification  
Time: 7 mins.  
Amplification: 6  
IC: tetanus serum + blood

Program: 430 liver detoxification  
930 lymph activation  
IC: blood  
I: flexible electrode on stomach and forehead  
O: modulation mat  
MD:chip

The patient told me after the first treatment that he felt even worse and the pressure in his head was more acute.

After the second treatment the pressure in his head had gone and he wanted to play football with his friends again. On the 4<sup>th</sup> day of treatment he explained that he had got a 'B' for maths and also felt much happier in general.

After testing following the 4th treatment no traces of vaccine stress were detected.

## Case 3: Child with acute ADS symptoms

A boy, 5 years old, came to me with his mother. While the mother explained to me what had brought her to me, the boy clambered all over the treatment couch in my consulting room and touched everything that was on my desk, talking constantly the whole time. He was so hyperactive that

finally he started doing somersaults on the carpet. His nursery had said that they couldn't do anything with the child.

### Testing

After testing with the BICOM (which required a supreme effort and no little patience on my part) I established the cause as vaccine damage and food intolerance.

My staff subsequently complained to me that this boy had been impossible, since during therapy he kept fidgeting and spitting and shouting out things like: you bitch, you dope, you ass-hole and worse.

### Therapy

Program: 191 detoxification  
Time: 5 mins.  
Amplification: 5  
IC: immune serum + blood

Program: 430 liver detoxification  
930 lymph activation  
IC: blood  
I: flexible electrode on stomach and forehead  
O: modulation mat  
MD:chip

After the first treatment the symptoms worsened. The mother was considering whether to continue. I talked to her and explained that these were reactions to the treatment. After the 2<sup>nd</sup> session my staff told me that the boy had been much more pleasant. His mother now wanted to continue treatment for her son. We treated him a total of 6 times and then the vaccine damage was gone and the child was much calmer and more sensible. I then successfully treated the food intolerances.

Today this little patient is completely different. At nursery there are no more complaints and a sense of calm has returned to the family. Despite successful treatment I recommended that the mother avoid all artificial food additives and fed the child natural and healthy products.

## CRY-BABIES

Cry-babies with acute sleep and development deficiencies: the parents do not get any peace at night and the child is roused from its sleep and starts crying. Parents cannot calm such children even when they get older. This is no longer a case of three-month colic. Often the only thing which helps is to take the child into your own bed. These

children often suffer from inexplicable fears, especially in the dark. They often sweat an unnatural amount at night and clearly have too much saliva even once they have stopped teething. Their limbs are constantly moving and such children cannot rest. Here is one such case.

## **Case 4: Cry-baby with acute development problems**

Child, 1 ½ years old, still cannot sit and crawl independently, not very aware of surroundings. The mother explained that her child had received a six-way inoculation at 8 weeks and became very ill as a result. She had a high fever and suffered fits. During this time, her parents did not get any rest at night. The child cried, was roused from her sleep and her mother found it difficult to calm her down again. The doctors told the mother that her child had autistic traits.

### Testing

I established by testing with BICOM that this child was showing signs of acute vaccine stress. She also reacted strongly to the Hib-tetanus-diphtheria-polio and whooping cough vaccine and less so to the measles-mumps jab.

### Therapy

Program: 191 detoxification

Time: 5 mins.

Amplification: 5

IC: immune serum + blood

I: flexible electrode on forehead

O: modulation mat

Liver, kidneys, lymphs and intestine were detoxified at the same time.

Program: 430 liver detoxification

930 lymph activation

IC: blood

I: electrode on stomach and forehead

O: modulation mat

Treatment once a week.

After 6 therapy sessions the child changed noticeably and began looking for eye contact and was more aware of her surroundings. After a further 6 treatments the child could sit by herself.

At the same time I tested and treated the child for food intolerances using BICOM. It was only now that the ergo therapy which the child had been

receiving for over a year began to work. I later treated the brain metabolism. Now, one year on this child can walk and is starting to speak simple words.

## **PATIENTS WITH HEADACHES, STOMACH ACHES, IMPAIRED VISION**

Often patients come to me with headaches, stomach aches and impaired vision. They often complain that they cannot concentrate. A few of them have a tremor in their hands which means that they cannot hold a full cup without spilling the contents. Some of them have great trouble trying to write. It is thought that tremors in particular are almost impossible to treat using conventional medicine. For complaints of this kind I always test for inoculations. In most cases I find a vaccine stress. Here is another particularly acute case.

## **Case 5: Man with tremor, migraine attacks, memory loss**

A man, 35 years old, suffers from a tremor in the left hand, migraine attacks and memory loss. One year previously he had a tetanus-hepatitis-B-malaria inoculation as a preventive measure before going to work abroad. The migraines started 4 weeks after the inoculation and the tremor developed within 4 to 5 months.

### Therapy after testing

Program: 191 detoxification

Time: 8 mins.

Amplification: 7

IC: immune serum + blood

Liver, kidneys, lymphs and intestine were detoxified at the same time.

Program: 430 liver detoxification

200 lymph acute-inflammatory

561 intestinal flora, to improve

IC: blood

For all programs mentioned:

I: flexible electrode on stomach and forehead

O: modulation mat

MD: chip

After 12 treatments the migraines and memory loss had disappeared and the tremor was as good as gone. Only in stressful situations did he report a slight temporary relapse.

## **PATIENTS WITH SKIN DISORDERS, NEURODERMATITIS, PSORIASIS**

For other patients with skin disorders such as neurodermatitis and psoriasis, especially children, I always test for inoculations. I frequently discover that whooping cough is acute in these patients. I recommend in these cases testing all stresses, such as moulds, heavy metals, food and especially inoculations since many immune serums contain mercury. In the package insert mercury appears under the heading thiomersal.

For these patients I use the following therapy.

### Therapy after testing an additional yeast stress

Program: 971 mycosis therapy  
561 intestinal flora, to improve  
IC: yeast fungi + spirulina + blood

Program: 191 detoxification  
IC: immune serums + blood  
Time and amplification after testing

I also treated using

Program: 530 metabolism therapy  
930 lymph activation  
IC: blood

For all programs mentioned:

I: flexible electrode on stomach and forehead  
O: modulation mat.

When this is cleared up I then treat the food intolerances.

## **PATIENTS WITH IMMUNE DEFICIENCIES**

I also test a large number of children with immune deficiencies for inoculations using the BICOM. Even in patients suffering from allergies, inoculations often play a major part. I would recommend that you always test patients with these disorders for vaccine stresses too. My experience has shown that at least 60 % of all allergy patients have a stress of this kind present.

## **PATIENTS WITH DIABETES**

Remember too that inoculations may be the cause of diabetes I + II. I have diagnosed a vaccine stress in many diabetics. With diabetes-II patients I have had good results using BICOM treatment. In con-

junction with a change in diet the blood sugar level soon returns to normal after the vaccine damage has been eliminated.

I have previously only treated a few diabetes-I patients for vaccine damage as I don't get many of these patients turning up at my practice since I cannot prescribe insulin. However, there are signs here too that this method works.

I urgently recommend you get yourself an inoculation test kit or immune serums to test out on your patients.

## **VACCINE DAMAGE – TOO HIGH A PRICE?**

Vaccine damage is much more widespread than conventional medicine will officially admit.

In my opinion, the key factor is the high number of inoculations. According to the pre-school inoculation schedule 33 inoculations are recommended whereas before 1970 there were only four. Those in favour of inoculations claim that thanks to them it has been possible to completely wipe out or significantly reduce certain diseases such as smallpox. However, there are also critics who should be taken seriously who reject this. Even if it is true, you must ask yourself whether this is not far too high a price to pay?

There is an urgent need to return to former inoculation practice which would considerably reduce the risks:

- Significant reduction in the number of inoculations (most important measure)
- Return to single inoculations instead of the multiple inoculations common today
- Inoculations at the earliest after the age of one
- In the case of premature births, inoculations at the earliest after the age of 3
- Only having the child inoculated when they are completely healthy.

The experiences which I have reported to you affect around 700 patients each year, of which:

- ca. 350 allergy patients
- ca. 200 ADS patients
- ca. 150 patients with other disorders.

I diagnose and treat all of these with BICOM, but not just for vaccine damage. The success rate is above 90 %, so after a relatively short time I acquired two further BICOM devices and now I am more than happy to recommend BICOM therapy to others.